

The assignment is due at the beginning of class on December 10, 2007.

It is quite obvious that statements and their connectives on the one hand, and sets and set connectives on the other hand behave somewhat analogously. The English mathematician GEORGE BOOLE (1815–1864) made this idea precise by describing what he called “algebra of logic”. Today we use the name “Boolean Algebra” in his honor instead. The axioms below were first formulated by the American mathematician EDWARD V. HUNTINGTON¹ (1874–1952):

A *Boolean Algebra* is a set \mathcal{B} together with two “connectives” \sqcap and \sqcup satisfying the following properties:

1. Closure Laws:

- (a) If A and B are two elements in \mathcal{B} , then $A \sqcap B$ is also an element in \mathcal{B} .
- (b) If A and B are two elements in \mathcal{B} , then $A \sqcup B$ is also an element in \mathcal{B} .

2. Commutative Laws:

- (a) $A \sqcap B = B \sqcap A$ for all elements A and B in \mathcal{B} .
- (b) $A \sqcup B = B \sqcup A$ for all elements A and B in \mathcal{B} .

3. Distributive Laws:

- (a) $A \sqcap (B \sqcup C) = (A \sqcap B) \sqcup (A \sqcap C)$ for all elements A, B and C in \mathcal{B} .
- (b) $A \sqcup (B \sqcap C) = (A \sqcup B) \sqcap (A \sqcup C)$ for all elements A, B and C in \mathcal{B} .

4. Identity Laws:

There are elements $N \in \mathcal{B}$ (called the *null element*) and $O \in \mathcal{B}$ (the *one element*) such that

- (a) $A \sqcap N = N$ and $A \sqcap O = A$ for all elements A in \mathcal{B} .
- (b) $A \sqcup O = O$ and $A \sqcup N = A$ for all elements A in \mathcal{B} .

5. Complement Law:

For every element A in \mathcal{B} there is an element B in \mathcal{B} such that $A \sqcap B = N$ and $A \sqcup B = O$.

6. Associative Laws:²

- (a) $A \sqcap (B \sqcap C) = (A \sqcap B) \sqcap C$ for all elements A, B and C in \mathcal{B} .
- (b) $A \sqcup (B \sqcup C) = (A \sqcup B) \sqcup C$ for all elements A, B and C in \mathcal{B} .

Let X be an arbitrary set. Then $\mathcal{P}(X)$ with the connectives \cap (in the role of \sqcap) and \cup (in the role of \sqcup) forms a Boolean Algebra.

¹Edward V. Huntington: *Sets of Independent Postulates for the Algebra of Logic*. Transactions of the American Mathematical Society 5 (1904), pp. 288-309.

²The Associative Laws can be deduced from the other five Boolean Algebra Laws.

Problem 1 (old) Let X be an arbitrary set. Let A and B be elements in $\mathcal{P}(X)$. Show the following

$$A \cap (A \cup B) = A.$$

(Similarly one obtains that $A \cup (A \cap B) = A$.)

Certain sets of statements with connectives \wedge (in the role of \cap) and \vee (in the role of \cup) also form Boolean Algebras.

What is meant by “certain” sets of statements? Our task at hand is to identify what sets of statements correspond to power sets.

Let us consider an example and start with one “generic” statement P . How many distinct propositional forms can we form involving this statement? A little bit of reflection will lead us on the following path: Every propositional form has a truth table, so the number of distinct propositional forms is limited by the number of distinct truth tables. Since a truth table involving the statement P has two rows, and since we have two choices for each row entry (T or F), there are at most 4 distinct truth tables, and therefore there are at most 4 distinct propositional forms. On the other hand it is easy to see that P , $\neg P$, $P \vee \neg P$ and $P \wedge \neg P$ are 4 distinct propositional forms contained in each Boolean Algebra containing P .

It is now boring to check that the following 4-element set indeed forms a Boolean Algebra:

$$\mathcal{S}_1 = \{P \wedge \neg P; P, \neg P; P \vee \neg P\}$$

\mathcal{S}_1 is called the “Boolean Algebra generated by the free statement P ”.

Problem 2 (old) Find the Boolean Algebra \mathcal{S}_2 generated by two free statements P and Q . How many elements does \mathcal{S}_2 have?

Problem 3 (old) Let A and B be elements in a Boolean Algebra \mathcal{B} . Show:

$$A \cap (A \sqcup B) = A.$$

Analogously one can obtain $A \sqcup (A \cap B) = A$.

*For a natural number n , let \mathcal{D}_n denote the set of the divisors of n . For example, $\mathcal{D}_{42} = \{1, 2, 3, 6, 7, 14, 21, 42\}$ and $\mathcal{D}_{12} = \{1, 2, 3, 4, 6, 12\}$. For $m, n \in \mathbb{N}$ let $m \cap n$ denote the greatest common divisor of n and m , and $m \sqcup n$ their least common multiple. For instance $6 \cap 4 = 2$ and $6 \sqcup 4 = 12$. It turns out that \mathcal{D}_{42} with these two operations \cap and \sqcup forms a Boolean Algebra, while \mathcal{D}_{12} does **not**.*

Problem 4 (old) Verify Boolean Algebra Laws 3, 4 and 5 for \mathcal{D}_{42} .

Problem 5 (old) 1. Show that \mathcal{D}_{12} does not form a Boolean Algebra.

2. Conjecture for which values of n the set \mathcal{D}_n forms a Boolean Algebra.

Problem 6 (old) Consider the relation “ \preceq ” on a Boolean Algebra \mathcal{B} defined by

$$A \preceq B \iff A \sqcup B = B$$

for $A, B \in \mathcal{B}$. Prove that \preceq is reflexive, anti-symmetric and transitive.

Problem 7 (old) Consider the Boolean Algebra \mathcal{S}_1 . Draw a *Hasse diagram* for \mathcal{S}_1 endowed with the partial order \preceq .

Problem 8 (old) Let \mathcal{B} be a Boolean Algebra with null-element N , partially ordered by \preceq . We say that $A \in \mathcal{B}$ is an **ATOM** of \mathcal{B} if N is the immediate predecessor of A .

1. Find all atoms of $\mathcal{P}(\{1, 2, 3, 4\})$.
2. Find all atoms of \mathcal{D}_{42} .
3. Assume additionally that \mathcal{B} has finitely many elements. Show that for every $B \in \mathcal{B}$ with $B \neq N$ there is an atom A such that $A \preceq B$.

Problem 9 (10 points) Find a Boolean Algebra with 8 elements that is a subset of $\mathcal{P}(\{1, 2, 3, 4\})$, but **not** the power set of a three-element subset of $\{1, 2, 3, 4\}$, then find its atoms and draw its Hasse diagram.

Problem 10 (10 points) Let A, B be two elements in a Boolean Algebra. Show the following:

1. $A \preceq B \Leftrightarrow A \sqcap B = A$.
2. If $A \sqcap B = N$ and $A \preceq B$, then $A = N$.

Problem 11 (10 points) Let A_1, A_2 be two atoms in a finite Boolean Algebra. Show the following:

1. The least upper bound of the set $\{A_1, A_2\}$ is the element $A_1 \sqcup A_2$.
2. If $A_1 \neq A_2$, then $A_1 \sqcap A_2 = N$.

Problem 12 (10 points) Given an element B in a finite Boolean Algebra \mathcal{B} , we let

$$\alpha(B) = \{A \in \mathcal{B} \mid A \preceq B \text{ and } A \text{ is an atom of } \mathcal{B}\}.$$

Let $A_1 \neq A_2$ be two atoms in \mathcal{B} . Show that $\alpha(A_1 \sqcup A_2) = \{A_1, A_2\}$.³

In the sequel, you may assume that results corresponding to those proved for two atoms in Problems 11 and 12 also hold for finitely many atoms.

Problem 13 (10 points) Let $B \neq N$ be an element in a finite Boolean Algebra \mathcal{B} , and suppose $\alpha(B) = \{A_1, A_2, A_3, \dots, A_k\}$ for some $k \in \mathbb{N}$ and some atoms $A_1, A_2, A_3, \dots, A_k$ of \mathcal{B} . Show:⁴

$$B = A_1 \sqcup A_2 \sqcup A_3 \sqcup \dots \sqcup A_k.$$

The next problem is the finite version of a general representation theorem for Boolean Algebras, proved by the American mathematician MARSHALL H. STONE⁵ (1903–1989):

³By Problem 8.3 we know that $\alpha(B) \neq \emptyset$ whenever $B \neq N$. Also note that $\alpha(N) = \emptyset$.

⁴*Hint:* Expect to use Boolean Algebra Law 5 along the way.

⁵Marshall H. Stone: *The Theory of Representation for Boolean Algebras*. Transactions of the American Mathematical Society 40 (1936), pp. 37-111.

Problem 14 (20 points) Let \mathcal{B} be a finite Boolean Algebra with k atoms for some $k \in \mathbb{N}$, and let \mathcal{A} denote the power set of the set of all atoms of \mathcal{B} .

1. Show that the function $\alpha : \mathcal{B} \rightarrow \mathcal{A}$, defined in Problem 12, is a bijection.
2. \mathcal{B} has 2^k elements.
3. Show that the identities $\alpha(B \sqcup B') = \alpha(B) \cup \alpha(B')$ and $\alpha(B \sqcap B') = \alpha(B) \cap \alpha(B')$ hold for all $B, B' \in \mathcal{B}$.