

Let  $\mathcal{K}$  denote the set of  $2 \times 2$  matrices of the form  $\begin{pmatrix} a & b \\ -b & a \end{pmatrix}$ , where  $a, b \in \mathbb{R}$ .

- a. Show: If  $A$  and  $B$  are elements in  $\mathcal{K}$ , then  $A + B \in \mathcal{K}$ .
- b. Show: If  $A$  and  $B$  are elements in  $\mathcal{K}$ , then  $A \cdot B \in \mathcal{K}$ .
- c. Show that  $A \cdot B = B \cdot A$  holds for all elements  $A, B \in \mathcal{K}$ .
- d. Show: If the matrix  $A = \begin{pmatrix} a & b \\ -b & a \end{pmatrix}$  in  $\mathcal{K}$  satisfies  $a^2 + b^2 \neq 0$ , then  $A$  has a matrix inverse. What is  $A^{-1}$ ? Does  $A^{-1}$  always lie in  $\mathcal{K}$ ? (Note that the identity matrix is an element of  $\mathcal{K}$ .)
- e. Show that we can identify  $\mathcal{K}$  with  $\mathbb{C}$ , i.e., find a bijection  $f : \mathcal{K} \rightarrow \mathbb{C}$  such that  $f(A + B) = f(A) + f(B)$  and  $f(A \cdot B) = f(A) \cdot f(B)$ . (This is easier than it sounds.)
- f. Under this identification, find the element  $I$  in  $\mathcal{K}$  that corresponds to  $i$  (there are actually two possible choices). Compute  $I \cdot I$ .
- g. Under this identification, what is the “conjugate” of an element in  $\mathcal{K}$ , what is the “modulus” of an element in  $\mathcal{K}$ ?